

An example of simple Lie superalgebra with several invariant bilinear forms

S.E.Konstein ^{*†}

I.E.Tamm Department of Theoretical Physics,
P. N. Lebedev Physical Institute,
117924, Leninsky Prospect 53, Moscow, Russia.

Abstract

Simple associative superalgebra with 2 independent supertraces is presented. Its commutant is a simple Lie superalgebra which has at least 2 independent invariant bilinear forms.

^{*}E-mail: konstein@lpi.ru

[†]This work was supported by the Russian Basic Research Foundation, grant 99-02-17916 and grant 00-15-96566.

1 Introduction

In this paper the associative superalgebra $\mathcal{A} = SH'_3(0)$ with two-dimensional space of supertraces is presented. It is shown that (i) it is simple, (ii) its commutant $\mathfrak{A}_1 = [SH'_3(0), SH'_3(0)]$ is a simple Lie superalgebra and (iii) $\mathfrak{A}_1$ has at least 2-dimensional space of nondegenerate bilinear invariant forms. Would $\mathfrak{A}_1$ be finitedimensional superalgebra having 2-dimensional space of invariant bilinear forms, it necessarily would not be simple.

The superalgebra $SH'_3(0)$ belongs to the class of *superalgebras of observables of N -particle Calogero model* which is denoted in [8] as $SH'_N(\nu)$, where ν is a coupling constant of this Calogero model [1].

Let $x_i \in \mathbb{R}$, $i = 1, \dots, N$.

The Hamiltonian of Calogero model after some similarity transformation attains the form

$$H_{Cal} = -\frac{1}{2} \sum_{i=1}^N \left[\frac{\partial^2}{\partial x_i^2} - x_i^2 + \nu \sum_{j \neq i} \frac{2}{x_i - x_j} \frac{\partial}{\partial x_i} \right] \quad (1)$$

which coincides with the operator

$$H = \frac{1}{2} \sum_{i=1}^N \{a_i^0, a_i^1\} \quad (2)$$

on the space of symmetric functions. The annihilation and creation operators a_i^α ($\alpha = 0, 1$ correspondingly) acting on the space of complex valued infinitely smooth functions of $x_1, x_2, \dots, x_N$ are expressed via the differential-difference Dunkl operators $D_i(x)$ [4]

$$a_i^\alpha = \frac{1}{\sqrt{2}}(x_i + (-1)^\alpha D_i(x)), \text{ where } D_i = \frac{\partial}{\partial x_i} + \nu \sum_{l \neq i} \frac{1}{x_i - x_l} (1 - K_{il}), \quad (3)$$

and $K_{ij} = K_{ji}$ are the operators of elementary transpositions

$$K_{ij}x_i = x_j K_{ij}, \quad K_{ij}x_k = x_k K_{ij} \text{ if } k \neq i \text{ and } k \neq j, \quad (4)$$

satisfying as a consequence the following relations:

$$K_{ij}a_i^\alpha = a_j^\alpha K_{ij}, \quad K_{ij}a_k^\alpha = a_k^\alpha K_{ij} \text{ if } k \neq i \text{ and } k \neq j, \quad K_{ij}^2 = 1. \quad (5)$$

The operators a_i^α satisfy the following commutation relations

$$[a_i^\alpha, a_j^\beta] = \epsilon^{\alpha\beta} \left(\delta_{ij} + \nu \delta_{ij} \sum_{l=1, l \neq i}^N K_{il} - \nu \delta_{i \neq j} K_{ij} \right),$$

where $\epsilon^{\alpha\beta} = -\epsilon^{\beta\alpha}$, $\epsilon^{01} = 1$, which lead [6] to the relations

$$[H, a_i^\alpha] = -(-1)^\alpha a_i^\alpha,$$

and, therefore, a_i^1 when acting on the Fock vacuum $|0\rangle$ such that $a_j^0|0\rangle = 0$ for any j , give all the wavefunctions of the model (2) and, consequently, of the model (1). In [7] Perelomov

operators [2], describing the wavefunctions of the model (1) are expressed via these operators a_i^α .

Definition 1. $SH_N(\nu)$ is the associative superalgebra of all polynomials in the operators K_{ij} and a_i^α with $\mathbb{Z}_2$ -grading π defined by the formula: $\pi(a_i^\alpha) = 1$, $\pi(K_{ij}) = 0$.

The superalgebra $SH_N(\nu)$ is called the algebra of observables.

It is clear that $SH_N(\nu) = SH_1(0) \otimes SH'_N(\nu)$, where $SH_1(0)$ is generated by the operators $a^\alpha = \sum_i a_i^\alpha / \sqrt{N}$ satisfying the commutation relations $[a^\alpha, a^\beta] = \epsilon^{\alpha\beta}$, while $SH'_N(\nu)$ is generated by the transpositions K_{ij} and by all linear combinations of the form $\sum_i \lambda_i a_i^\alpha$ with coefficients λ_i satisfying the condition $\sum_i \lambda_i = 0$.

The group algebra $\mathbb{C}[S_N]$ of the permutation group S_N , generated by elementary transpositions K_{ij} is a subalgebra of $SH'_N(\nu)$.

Definition 2. The supertrace on an arbitrary associative superalgebra $\mathcal{A}$ is complex-valued linear function $str(\cdot)$ on $\mathcal{A}$ which satisfies the condition

$$str(fg) = (-1)^{\pi(f)\pi(g)} str(gf) \quad (6)$$

for every $f, g \in \mathcal{A}$ with definite parity.

It is proven in [8] that $SH_N(\nu)$ has nontrivial supertraces and the dimension of the space of the supertraces on $SH_N(\nu)$ is equal to the number of partitions of N into the sum of positive odd integers.¹ In particular, $SH_2(\nu)$ has only one supertrace and $SH_3(\nu)$ has two independent supertraces. Since $SH_N(\nu) = SH_1(0) \otimes SH'_N(\nu)$ and $SH_1(0)$ has one supertrace, it follows that the numbers of supertraces on $SH_N(\nu)$ and on $SH'_N(\nu)$ are the same.

Every supertrace $str(\cdot)$ defines the bilinear form B_{str} :

$$B_{str}(f, g) = str(fg). \quad (7)$$

Since the complete set of null-vectors of every invariant bilinear form constitutes the two-sided ideal of the algebra, the supertrace is a good tool for finding such ideals.

It should be noticed that the existence of supertraces depends on the choice of $\mathbb{Z}_2$ -grading π , while the existence of an ideals does not depend on the grading.

In the well known case of $SH'_2(\nu)$ corresponding to the usual two-particle Calogero model the only supertrace defines the bilinear form (7) which degenerates when ν is half-integer [5], hence, the superalgebra $SH'_2(\nu)$ has an ideal for these values of ν . When $N \geq 3$ the situation is more complicated as $SH'_N(\nu)$ has more than one supertrace. In the case of finitedimensional superalgebra this would be sufficient for existence of ideals but for infinitedimensional superalgebras under consideration the existence of ideals is not completely investigated yet.

It is shown in [10] that in the superalgebras $SH'_3(\nu)$ with $\nu = n \pm 1/3$ or $\nu = n + 1/2$ (n is an integer) there exists one supertrace with degenerated corresponding bilinear form (7) and that for the other values of ν all the bilinear forms corresponding to supertraces are non-degenerate.

One can suggest that for the latest case the corresponding superalgebras are simple.

Below it is shown that the associative superalgebra $SH'_3(0)$ is simple and that its commutant is simple Lie superalgebra.

¹The analogous result for Calogero model based on arbitrary root system [3] is presented in [11].

The paper is organized as follows. In Section 2 the generating elements of associative superalgebra $SH'_3(0)$ are presented together with the relations between them. The simplicity of $SH'_3(0)$ is proved in Section 3 and in the last section the simplicity of $[SH'_3(0), SH'_3(0)]$ is established.

2 The algebra $SH'_3(0)$

In this Section we present the generating elements of associative superalgebra $\mathcal{A} = SH'_3(0)$ and relations between them.

Let $\lambda = \exp(2\pi i/3)$. As a basis in $\mathbb{C}[S_3]$ let us choose the vectors

$$\begin{aligned} L_k &= \frac{1}{3}(\lambda^k K_{12} + K_{23} + \lambda^{-k} K_{31}), \\ Q_k &= \frac{1}{3}(1 + \lambda^k K_{12} K_{13} + \lambda^{-k} K_{12} K_{23}), \\ L_{i+3} &= L_i, \quad Q_{i+3} = Q_i, \\ L_{\pm} &\stackrel{def}{=} L_{\pm 1}, \quad Q_{\pm} \stackrel{def}{=} Q_{\pm 1}. \end{aligned}$$

Instead of the generating elements a_i^α let us introduce the vectors

$$\begin{aligned} x^\alpha &= a_1^\alpha + \lambda a_2^\alpha + \lambda^2 a_3^\alpha, \quad x \stackrel{def}{=} x^0, \quad x^+ \stackrel{def}{=} x^1, \\ y^\alpha &= a_1^\alpha + \lambda^2 a_2^\alpha + \lambda a_3^\alpha, \quad y \stackrel{def}{=} y^0, \quad y^+ \stackrel{def}{=} y^1, \end{aligned}$$

which are derived from a_i^α by subtracting the center of mass $a^\alpha = \frac{1}{3} \sum_{i=1,2,3} a_i^\alpha$.

The Lie algebra $\mathfrak{sl}_2$ of inner automorphisms of $SH'_3(0)$ is generated by the generators $T^{\alpha\beta}$

$$T^{\alpha\beta} = \frac{1}{3}(x^\alpha y^\beta + x^\beta y^\alpha). \quad (8)$$

These generators satisfy the usual commutation relations

$$[T^{\alpha\beta}, T^{\gamma\delta}] = \epsilon^{\alpha\gamma} T^{\beta\delta} + \epsilon^{\alpha\delta} T^{\beta\gamma} + \epsilon^{\beta\gamma} T^{\alpha\delta} + \epsilon^{\beta\delta} T^{\alpha\gamma}, \quad (9)$$

and act on generating elements x^α and y^α as follows:

$$[T^{\alpha\beta}, x^\gamma] = \epsilon^{\alpha\gamma} x^\beta + \epsilon^{\beta\gamma} x^\alpha, \quad [T^{\alpha\beta}, y^\gamma] = \epsilon^{\alpha\gamma} y^\beta + \epsilon^{\beta\gamma} y^\alpha \quad (10)$$

leaving the group algebra $\mathbb{C}[S_3]$ invariant: $[T^{\alpha\beta}, K_{ij}] = 0$.

Clearly, $SH'_3(0)$ decomposes into the infinite direct sum of finitedimensional irreducible representations of this $\mathfrak{sl}_2$. Let $\mathcal{A}_s$ (s is a spin) be the direct sum of all $2s + 1$ -dimensional irreducible representations in this decomposition. So

$$\mathcal{A} = \mathcal{A}_0 \oplus \mathcal{A}_{1/2} \oplus \mathcal{A}_1 \oplus \dots \quad (11)$$

Evidently, $\mathcal{A}_0$ is a subalgebra of $\mathcal{A} = SH'_3(0)$.

As every element of the subspace $\mathcal{A}_s$ for $s > 0$ is a linear combinations of the vectors of the form $[f, T^{\alpha\beta}]$ where $T^{\alpha\beta}$ is defined in (8), every supertrace vanishes on this subspace and has non-trivial values only on the associative subalgebra $\mathcal{A}_0 \subset SH'_3(0)$ of all $\mathfrak{sl}_2$ -singlets.

Let $m = \frac{1}{4}\{x^\alpha, y_\alpha\}$. Clearly, m is a singlet under the action of $\mathfrak{sl}_2$ (8); m can also be expressed in the form

$$m = \frac{1}{2}(x^\alpha y_\alpha + 3) \text{ or, equivalently,} \quad (12)$$

$$m = \frac{1}{2}(y_\alpha x^\alpha - 3). \quad (13)$$

In these formulas the greek indices are lowered and raised with the help of the antisymmetric tensor $\varepsilon^{\alpha\beta}$: $a^\alpha = \sum_\beta \varepsilon^{\alpha\beta} a_\beta$.

The generating elements $x^\alpha, y^\alpha, Q_i, L_i$ and the element m satisfy the following relations:

$$L_i L_j = \delta_{i+j} Q_j, \quad L_i Q_j = \delta_{i-j} L_j, \quad (14)$$

$$Q_i L_j = \delta_{i+j} L_j, \quad Q_i Q_j = \delta_{i-j} Q_j, \quad (15)$$

where $\delta_0 = 1$ and $\delta_i = 0$ if $i \neq 0$,

$$L_i x^\alpha = y^\alpha L_{i+1}, \quad L_i y^\alpha = x^\alpha L_{i-1}, \quad (16)$$

$$Q_i x^\alpha = x^\alpha Q_{i+1}, \quad Q_i y^\alpha = y^\alpha Q_{i-1}, \quad (17)$$

$$L_i m = -m L_i, \quad Q_i m = m Q_i, \quad (18)$$

$$\begin{aligned} [x, x^+] &= 0, & [y, y^+] &= 0, & [y, x^+] &= [x, y^+] = 3, \\ & & & & [x, y] &= [x^+, y^+] = 0, \end{aligned} \quad (19)$$

and

$$\begin{aligned} [m, x^\alpha] &= \frac{3}{2}(x^\alpha), \\ [m, y^\alpha] &= -\frac{3}{2}(y^\alpha), \end{aligned} \quad (20)$$

Corollary 1. There exists $\mathbb{Z}_3$ grading ρ on $SH'_3(0)$ defined by the formulas:

$$\begin{aligned} \rho(Q_i) &= 0, \\ \rho(L_i) &= -i, \\ \rho(x^\alpha) &= 1, \\ \rho(y^\alpha) &= -1, \end{aligned}$$

such that $fQ_i = Q_i f$ for any element $f \in SH'_3(0)$ with $\rho(f) = 0$. Particularly, $[(x^\alpha)^3, Q_i] = [(y^\alpha)^3, Q_i] = 0$.

Corollary 2. There exists the antiautomorphism τ of associative algebra $SH'_3(0)$:

$$\begin{aligned} \tau(x^\alpha) &= y^{1-\alpha}, \quad \tau(y^\alpha) = x^{1-\alpha}, \quad \tau(m) = m, \\ \tau(L_i) &= L_{-i}, \quad \tau(Q_i) = Q_i. \end{aligned}$$

Every supertrace on $SH'_3(\nu)$ is completely determined by its values on $\mathbb{C}[S_3] \subset SH'_3(\nu)$, i.e., by the values $str(1)$, $str(K_{12}) = str(K_{23}) = str(K_{31})$ and $str(K_{12}K_{23}) = str(K_{12}K_{13})$,

which are consistent with (6) and admit the extension of the supertrace from $\mathbb{C}[S_3]$ to $SH'_3(\nu)$ in a unique way, if and only if $str(K_{ij}) = \nu(-2str(1) - str(K_{12}K_{23}))$ [8].

It was noticed above that only trivial representations of $\mathfrak{sl}_2$ can contribute to any supertrace on $SH'_3(0)$. Evidently, the associative algebra $\mathcal{A}_0$ of $\mathfrak{sl}_2$ -singlets contains only the polynomials of m with coefficients in $\mathbb{C}[S_3]$.

To describe the restrictions of all the supertraces on $\mathcal{A}_0$, it is convenient to use the generating functions, which are computed in [10] for any values of ν , and for $\nu = 0$ take the following form:

$$\mathcal{L}_i \stackrel{def}{=} str(e^{2/3\xi m} L_i) = 0, \quad (21)$$

$$\mathcal{Q}_i \stackrel{def}{=} str(e^{2/3\xi m} Q_i) = \frac{P_i}{\Delta}, \quad (22)$$

where

$$P_0 = -2S_1 + \frac{S_2}{2} (e^{2\xi} + e^{-2\xi} - 2e^\xi - 2e^{-\xi}), \quad (23)$$

$$P_+ = \frac{2}{3}S_1 (-e^{2\xi} + 2e^{-\xi}) + \frac{S_2}{2} (e^{-2\xi} - 2e^\xi + 3), \quad (24)$$

$$P_- = \frac{2}{3}S_1 (-e^{-2\xi} + 2e^\xi) + \frac{S_2}{2} (e^{2\xi} - 2e^{-\xi} + 3) \quad (25)$$

$$\Delta = exp(-3\xi)(exp(3\xi) + 1)^2. \quad (26)$$

Here S_1, S_2 are arbitrary parameters, specifying the supertrace in the two-dimensional space of supertraces:

$$S_1 = -2str(1) - str(K_{12}K_{23}), \quad S_2 = \frac{8}{3}str(1) - \frac{2}{3}str(K_{12}K_{23}).$$

3 The simplicity of $SH'_3(0)$.

To prove the simplicity of $SH'_3(0)$ it is sufficient to prove that elements Q_i belong to any nonzero ideal $\mathcal{I}$.

Let $f \in \mathcal{I}$, $f = f_0 + f_{1/2} + f_1 + \dots$, where $f_s \in \mathcal{A}_s$. As $[T^{\alpha\beta}, f] \in \mathcal{I}$ then $f_s \in \mathcal{I}$. So each ideal in $\mathcal{A}$ is graded.

Proposition 1. *Let $\mathcal{I} \subset SH'_3(0)$ be two-sided nonzero ideal. Then there exist nonzero polynomial $f(x^\alpha, y^\alpha)$ and a polynomial $g(x^\alpha, y^\alpha)$ such that $fQ_0 + gL_0 \in \mathcal{I}$.*

Indeed, let $r \in \mathcal{I}$, $r \neq 0$. Then at least one of the polynomials $rQ_0, rL_0, rxQ_0, rxL_0, ryQ_0, ryL_0$ has the desired form.

Proposition 2. *Let $\mathcal{I} \subset SH'_3(0)$ be two-sided nonzero ideal. Then there exist nonzero polynomial $f_0(m)$ and a polynomial $g_0(m)$ such that $f_0Q_0 + g_0L_0 \in \mathcal{I}$.*

Let $f(x^\alpha, y^\alpha) \neq 0$ and $h_0 = fQ_0 + gL_0 \in \mathcal{I} \cap \mathcal{A}_s$ with some polynomial $g(x^\alpha, y^\alpha)$. Let

$$h_{k+1} = [T^{00}, h_k] \quad (27)$$

Evidently, $h_k \in \mathcal{I}$ for any k . Then there exists such integer K that $h_K \neq 0$ and $h_{K+1} = 0$. The vector h_K is the highest vector of some irreducible representation of $\mathfrak{sl}_2$ algebra (8) and

has the form

$$h_K = \sum_{i=0}^{2s} \left((x)^i (y)^{2s-i} q_i(m) Q_0 + (x)^i (y)^{2s-i} p_i(m) L_0 \right) \quad (28)$$

where some polynomial $q_n(m) \neq 0$. Then $F = (y^+)^n (x^+)^{2s-n} h_K \in \mathcal{I}$. Let F_0 be the singlet part of F . Evidently, $F_0 \neq 0$, $F_0 \in \mathcal{I}$, and $F_0 = f_0(m) Q_0 + g_0(m) L_0$ with nonzero polynomial $f_0(m)$.

Proposition 3. *Let $\mathcal{I} \subset SH'_3(0)$ be two-sided nonzero ideal. Then there exists nonzero polynomial $f_0(m)$ such that $F_0 = f_0(m) Q_0 \in \mathcal{I}$.*

Let $F = f(m) Q_0 + g(m) L_0 \in \mathcal{I}$, $f(m) \neq 0$. Then $mf(m) Q_0 = \frac{1}{2}(mF + Fm) \in \mathcal{I}$ due to commutational relations.

Proposition 4. *Let $\mathcal{I} \subset SH'_3(0)$ be two-sided nonzero ideal. Then there exist nonzero polynomials $f_i(m)$ such that $F_i = f_i(m) Q_i \in \mathcal{I}$ for $i = 0, \pm 1$.*

Let $F_0 = f_0(m) Q_0 \in \mathcal{I}$ is nonzero polynomial from Proposition 3. Let us denote the singlet part of arbitrary element $F \in \mathcal{A}$ as $(F)_0$. Then $F_1 = (y^+ F_0 x)_0 = f_1(m) Q_1$ and $F_2 = (x^+ F_0 y)_0 = f_2(m) Q_2$ are nonzero and belong to $\mathcal{I}$.

Proposition 5. *Let $\mathcal{I}_i = \{f(m) \in \mathbb{C}[m] : f(m) Q_i \in \mathcal{I}\}$. Then $\mathcal{I}_i$ are nonzero ideals in $\mathbb{C}[m]$.*

This proposition follows trivially from Proposition 4.

Hence there exist polynomials $\varphi_i(m) = m^{k_i} + \dots$ such that $\mathcal{I}_i = \varphi_i(m) \cdot \mathbb{C}[m]$.

Proposition 6. *The polynomials $\varphi_i(m)$ generating the ideals $\mathcal{I}_i$ satisfy the relations*

$$\varphi_0(m) = \pm \varphi_0(-m), \quad (29)$$

$$\varphi_1(m) = \pm \varphi_2(-m). \quad (30)$$

Indeed, $\varphi_0(-m) Q_0 = L_0 \varphi_0(m) Q_0 L_0 \in \mathcal{I}$, hence $\varphi_0(-m) \in \mathcal{I}_0$ and all the roots of $\varphi_0(m)$ are the roots of $\varphi_0(-m)$. Further, $L_1 \varphi_1(m) Q_1 L_2 = \varphi_1(-m) Q_2$ and, as a consequence, $\varphi_1(-m) = p(m) \varphi_2(m)$ with some nonzero polynomial $p(m)$. Analogously, $\varphi_2(-m) = q(m) \varphi_1(m)$ with some nonzero polynomial $q(m)$. So $p(m) q(-m) = 1$, and $\varphi_2(-m) = \pm \varphi_1(m)$.

Now we can prove the following theorem:

Theorem 1. *The associative superalgebra $\mathcal{A} = SH'_3(0)$ is simple.*

We will prove that arbitrary nonzero ideal in $\mathcal{A}$ contains the element $1 \in \mathcal{A}$. Let $\mathcal{I}$ be nonzero ideal in $\mathcal{A}$ and $\varphi_i(m)$ generate the ideals $\mathcal{I}_i$ described above. So $\varphi_i(m) Q_i \in \mathcal{I}$ and hence $y^3 \varphi_i(m) Q_i (x^+)^3 \in \mathcal{I}$. Consider the $\mathfrak{sl}_2$ -singlet part of this element which also belongs to $\mathcal{I}$ and can be computed with the help of (20):

$$(y^3 \varphi_i(m) Q_i (x^+)^3)_0 = P(m) \varphi_i(m + 9/2) Q_i \in \mathcal{I}, \quad (31)$$

$$\text{where } P(m) = \text{const}(m + 3/2)(m + 3)(m + 9/2) \neq 0. \quad (32)$$

So $P(m) \varphi_i(m + 9/2) \in \mathcal{I}_i$ and

$$P(m) \varphi_i(m + 9/2) = p_i(m) \varphi_i(m)$$

with some polynomials $p_i(m)$. Since $\varphi_0(m)$ is even or odd polynomial, the real part of the rightest root of $\varphi_0(m)$ is non negative, and so this rightest root is not a root of polynomial

$(m + 3/2)(m + 3)(m + 9/2)\varphi_0(m + 3/2)$. So $\varphi_0(m)$ has no roots and $\varphi_0(m) = 1$. One can apply the same consideration to the function $\varphi(m) = \varphi_1(m)\varphi_2(m)$ which is even due to Proposition 6 and satisfies the equation

$$P(m)^2\varphi(m + 9/2) = p_1(m)p_2(m)\varphi(m).$$

So $\varphi_1(m)\varphi_2(m) = 1$ and $\varphi_1(m) = \varphi_2(m) = 1$.

In such a way, $Q_i \in \mathcal{I}$ and $1 = Q_0 + Q_1 + Q_2 \in \mathcal{I}$.

4 Lie superalgebra $\mathcal{A}^L$

Now let us consider Lie superalgebra $\mathfrak{A} = \mathcal{A}^L$ which consists of elements of associative algebra $\mathcal{A} = SH'_3(0)$ with operation $[f, g] = fg - (-1)^{\pi(f)\pi(g)}gf$. This Lie superalgebra has an ideal $\mathfrak{A}_1 = [\mathfrak{A}, \mathfrak{A}]$ with the following evident properties

- (i) $\text{codim}\mathfrak{A}_1 = 2$,
- (ii) $\mathcal{A}_{1/2} \oplus \mathcal{A}_1 \oplus \dots \subset \mathfrak{A}_1$.

Let Z be the center of $SH'_3(0)$. Obviously, $Z \subset \mathcal{A}_0$ and so every element of Z is a polynomial of m with coefficients in $\mathbb{C}[S_3]$.

Proposition 7. $Z = \mathbb{C}$.

Let $f(m) \in Z$. As m is even then f is even element also. As $mL_i = -L_im$, the relation $[f, m] = 0$ gives that $f(m) = \sum f_i(m)Q_i$. Due to equalities $[(x)^3, Q_i] = 0$ and (20) one has $[(x)^3, f(m)] = (x)^3(f(m) - f(m - 9/2))$ and as a consequence $f(m) = f(m - 9/2)$. So $f(m)$ does not depend on m , $f(m) \in \mathbb{C}[S_3]$. To finish the proof it is sufficient to commute f with x or y .

Proposition 8. $Z \cap \mathfrak{A}_1 = \{0\}$.

As there exists such supertrace that $\text{str}(1) \neq 0$ and $Z = \mathbb{C}$, it follows from $\text{str}(Z \cap \mathfrak{A}_1) = 0$ that $Z \cap \mathfrak{A}_1 = \{0\}$.

Proposition 9. Let $\mathcal{I} \subset \mathfrak{A}_1$ be an ideal and $\mathcal{I}_{\text{even}} = \{0\}$. Then $\mathcal{I}_{\text{odd}} = \{0\}$.

Let $r \in \mathcal{I}_{\text{odd}}$. Due to $\mathfrak{sl}_2$ invariance one can choose r in the form

$$r = \sum_{i=0,1,2} \sum_{n=0}^{2s} \left((x)^n (y)^{2s-n} f_{ni}(m) Q_i + (x)^n (y)^{2s-n} g_{ni}(m) L_i \right)$$

with odd value of $2s$. As $(x)^3 \in \mathcal{A}_{3/2} \subset \mathfrak{A}_1$ then $[(x)^3, r] \in \mathcal{I}_{\text{even}}$ and $\{(x)^3, r\} = 0$. So the following relations take place:

$$\sum_{i=0,1,2} \sum_{n=0}^{2s} (x)^{n+3} (y)^{2s-n} (f_{ni}(m) + f_{ni}(m + 9/2)) Q_i = 0 \quad (33)$$

and so $f_{ni} = 0$.

Further, it follows from $0 = \{(x)^3, r\} \simeq \sum_{i=0,1,2} \sum_{n=0}^{2s} ((x)^3 + (y)^3) (x)^n (y)^{2s-n} g_{ni}(m) L_i$ that $g_{ni}(m) = 0$ also. Here the sign $\simeq$ is used to denote the equality up to polynomials of lesser degrees.

Theorem 2 Lie superalgebra $\mathfrak{A}_1$ is simple.

This Theorem follows from the next two general theorems proven by S.Montgomery in [9].

Theorem 3 Let $\mathcal{A}$ be associative simple superalgebra and $\mathcal{I} \subset [\mathcal{A}^L, \mathcal{A}^L]$ be a graded ideal of $[\mathcal{A}^L, \mathcal{A}^L]$ such that $\mathcal{I} \neq [\mathcal{A}^L, \mathcal{A}^L]$. Then $\mathcal{I}_3 = \{0\}$, where $\mathcal{I}_1 = [\mathcal{I}, \mathcal{I}]$, $\mathcal{I}_2 = [\mathcal{I}_1, \mathcal{I}_1]$, $\mathcal{I}_3 = [\mathcal{I}_2, \mathcal{I}_2]$.

Theorem 4. Let $\mathcal{A}$ be associative simple superalgebra and $\mathcal{I} \subset [\mathcal{A}^L, \mathcal{A}^L]$ be a graded ideal of $[\mathcal{A}^L, \mathcal{A}^L]$ such that $[\mathcal{I}_{even}, \mathcal{I}] \subseteq Z(\mathcal{A})$ where $Z(\mathcal{A})$ is the center of $\mathcal{A}$. Then $\mathcal{I}_{even} \subseteq Z(\mathcal{A})$.

Indeed, let $\mathcal{I} \subset \mathfrak{A}_1$ be an ideal such that $\mathcal{I} \neq \mathfrak{A}_1$. Due to Theorem 3 one can consider that $[\mathcal{I}, \mathcal{I}] = \{0\}$. Then it follows from Theorem 4 and Proposition 8 that $\mathcal{I}_{even} = \{0\}$. Further, Proposition 9 gives that $\mathcal{I}_{odd} = \{0\}$, and so $\mathcal{I} = \{0\}$.

Theorem 5 Simple Lie superalgebra $\mathfrak{A}_1$ has at least 2 independent bilinear invariant forms.

Two-dimensional space of supertraces $str(\cdot)$ on $\mathcal{A} = SH'_3(0)$ generates some space of invariant bilinear forms $B(u, v) = str(uv)$ on $\mathfrak{A}_1$. This space is also 2-dimensional. Indeed, let some supertrace $str_0(\cdot)$ on $\mathcal{A}$ leads to bilinear form B_0 on $\mathfrak{A}_1$ which is equal to zero identically. The elements x, y^+, xQ_1 and y^+Q_1 belong to $\mathcal{A}_{1/2} \subset \mathfrak{A}_1$. So, $0 = B_0(x, y^+) - B_0(y^+, x) = str_0([x, y^+]) = 3str_0(1)$ and $0 = B_0(x, y^+Q_1) - B_0(y^+, xQ_1) = str_0([x, y^+]Q_1) = 3str_0(Q_1)$. It follows from (22) that $str_0(1) = -\frac{1}{6}(S_1 - \frac{3}{2}S_2)$ and $str_0(Q_1) = \frac{1}{6}(S_1 + \frac{3}{2}S_2)$. So $S_1 = S_2 = 0$ and str_0 is equal to zero identically.

Acknowledgments

The author is very grateful to M. Vasiliev and to D. Leites for useful discussions.

References

- [1] F. Calogero, J. Math. Phys., **10** (1969) 2191, 2197; *ibid*, **12** (1971) 419.
- [2] A. M. Perelomov, Teor. Mat. Fiz., **6** (1971) 364.
- [3] M. A. Olshanetsky and A. M. Perelomov, Phys. Rep., **94** (1983) 313.
- [4] C.F. Dunkl, Trans. Am. Math. Soc., **311** (1989) 167.
- [5] M.A. Vasiliev, JETP Letters, **50** (1989) 344; Int. J. Mod. Phys., **A6** (1991) 1115.
- [6] L. Brink, H. Hansson and M.A. Vasiliev, Phys. Lett., **B286** (1992) 109.
- [7] L. Brink, H. Hansson, S.E. Konstein and M.A. Vasiliev, Nucl. Phys., **B401** (1993) 591.
- [8] S.E. Konstein and M.A. Vasiliev, J. Math. Phys., **37** (1996) 2872.
- [9] S. Montgomery, J. of Algebra, **195** (1997) 558.
- [10] S.E. Konstein, Theor. Math. Phys., **116** (1998); hep-th/9803213.
- [11] S.E. Konstein, math-ph/9904032.